

11-28-05;03:44PM;

;1-732-321-3030

4/ 14

NOV 28 2005

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICATION NO. 10/736,087	FILING DATE 12/15/2003	FIRST NAMED INVENTOR Joseph C. Walsh	ATTORNEY DOCKET NO. 2003P88074US
-------------------------------	---------------------------	---	-------------------------------------


Response To OFFICIAL ACTION

EXAMINER Jones, Dameron Levest	
ART UNIT 1618	PAGE NUMBER 2

AMENDMENTS TO THE CLAIMS


In the Claims, please make the following amendments:

1. (Withdrawn) A method of preparing an ^{18}F -FLT precursor having the following formula:




wherein X is t-butoxycarbonyl, methyl, methoxymethyl ethyl, isobutyl, tetrahydropyranyl ether, tetrahydrofuranyl ether, methoxymethyl ether, bis-(2-chloroethoxy)methyl ether, 1-ethoxyethyl ether, or 1-methyl-1-methoxyethyl ether; P is an amine protecting group; and L is a leaving group, comprising the steps of:

- a. reacting a compound having the following formula:



with a reagent that opens the 2,3'-anhydro ring to produce 3'-beta-hydroxy thymidine;

- b. protecting the 5'hydroxyl group to produce a compound having the following formula:



wherein X is the same as defined above;

- c. incorporating a leaving group at the 3'-position to produce a compound having the following formula:


IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.
10/736,087	12/15/2003	Joseph C. Walsh	2003P88074US

Response To OFFICIAL ACTION

EXAMINER
Jones, Dameron Levest

ART UNIT	PAGE NUMBER
1618	3



wherein X and L are the same as defined above; and


- d. protecting the 3-N amine moiety to produce the precursor.
2. (Withdrawn) The method according to Claim 1, wherein the amine protecting group is selected from the group consisting of tert-butoxycarbamate, isopropyl carbamate, pivaloyloxymethyl carbamate, methyl carbamate, allyl carbamate, N-tetrahydropyran, N-tetrahydrofuran; t-butylamide, and N-pyrrolidinomethylamide.
3. (Withdrawn) The method according to Claim 1, wherein the amine protecting group is t-butoxycarbonyl.
4. (Withdrawn) The method according to Claim 1, wherein X is t-butoxycarbonyl, methyl, methoxymethyl ethyl, or isobutyl.
5. (Withdrawn) The method according to Claim 1, wherein X is t-butoxycarbonyl.
6. (Withdrawn) The method according to Claim 1, wherein L is selected from the group consisting of benzenesulfonyl, methylsulfonyl, p-methylbenzenesulfonyl, 4-nitrobenzene sulfonyl, p-bromobenzenesulfonyl, trifluoromethylsulfonyl, trichloroacetimidate, 2,2,2-trifluoroethanesulfonyl, and imidazolesulfonyl.
7. (Withdrawn) The method according to Claim 1, wherein L is nosylate.
8. (Withdrawn) The method according to Claim 1, wherein the precursor is 5'-O-Boc-3'-β-nosyl-3-N-Boc thymidine.
9. (Withdrawn) The method according to Claim 1, wherein the 2,3'-anhydro ring is opened by reacting 2,3'-anhydrothymidine with NaOH, KOH, LiOH, or tetrabutylammonium hydroxide.
10. (Withdrawn) A method of preparing an ¹⁸F-FLT precursor having the following formula:

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICATION NO. 10/736,087	FILING DATE 12/15/2003	FIRST NAMED INVENTOR Joseph C. Walsh	ATTORNEY DOCKET NO. 2003P88074US
-------------------------------	---------------------------	---	-------------------------------------


Response To OFFICIAL ACTION

EXAMINER	
Jones, Dameron Levest	
ART UNIT	PAGE NUMBER
1618	4




wherein P is an amine protecting group and L is a leaving group, comprising the steps of:

- a. reacting a compound having the following formula:




with a reagent that opens the 2,3'-anhydro ring to produce 3'-beta-hydroxy thymidine;

- b. reacting the reaction product of step (a) with BOC_2O to produce a compound having the following formula:



- c. incorporating a leaving group at the 3'-position to produce a compound having the following formula:



wherein L is the same as defined above; and

- d. protecting the 3-N amine moiety to produce the precursor.

11. (Withdrawn) The method according to Claim 10, wherein the amine protecting group is selected from the group consisting of tert-butoxycarbamate, isopropyl

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.
10/738,087	12/15/2003	Joseph C. Walsh	2003P88074US

Response To OFFICIAL ACTION

EXAMINER	
Jones, Dameron Levest	
ART UNIT	PAGE NUMBER
1618	5

carbamate, pivaloyloxymethyl carbamate, methyl carbamate, allyl carbamate, *N*-tetrahydropyran, *N*-tetrahydrofuran; *t*-butylamide, and *N*-pyrrolidinomethylamide.

12. (Withdrawn) The method according to Claim 10, wherein the amine protecting group is *t*-butoxycarbonyl.

13. (Withdrawn) The method according to Claim 10, wherein L is selected from the group consisting of benzenesulfonyl, methylsulfonyl, *p*-methylbenzenesulfonyl, 4-nitrobenzene sulfonyl, *p*-bromobenzenesulfonyl, trifluoromethylsulfonyl, trichloroacetimidate, 2,2,2-trifluoroethanesulfonyl, and imidazolesulfonyl.

14. (Withdrawn) The method according to Claim 10, wherein L is nosylate.

15. (Withdrawn) The method according to Claim 10, wherein the precursor is 5'-*O*-Boc-3'- β -nosyl-2-*N*-Boc thymidine.

16. (Withdrawn) The method according to Claim 10, wherein the 2,3'-anhydro ring is opened by reacting 2,3'-anhydrothymidine with NaOH, KOH, LiOH, alkylammonium hydroxides such as tetrabutylammonium hydroxide.

17. (Withdrawn) A method for preparing an ^{18}F -FLT precursor comprising:

- converting thymidine into 2,3'-anhydrothymidine;
- opening the 2,3'-anhydro ring to produce 3'-beta-hydroxy thymidine;
- protecting the 5'-hydroxy with *t*-butoxycarbonyl;
- incorporating a leaving group at the 3'-position; and
- protecting the 3-N amine to produce the precursor.

18. (Withdrawn) A method according to Claim 17, wherein the 2,3'-anhydro ring is opened by reacting 2,3'-anhydrothymidine with NaOH, KOH, LiOH, or tetrabutylammonium hydroxide.

19. (Withdrawn) A method according to Claim 17, wherein the 5'-hydroxy is protected by reacting the reaction product of step (b) with BOC_2O .

20. (Withdrawn) A method according to Claim 17, wherein the leaving group is a sulfonate ester.


IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.
10/736,087	12/15/2003	Joseph C. Walsh	2003P88074US

Response To OFFICIAL ACTION

EXAMINER	
Jones, Dameron Levest	
ART UNIT	PAGE NUMBER
1618	6

21. (Withdrawn) A method according to Claim 17, wherein L is selected from the group consisting of benzenesulfonyl, methylsulfonyl, p-methylbenzenesulfonyl, 4-nitrobenzene sulfonyl, p-bromobenzenesulfonyl, trifluoromethylsulfonyl, trichloroacetimidate, 2,2,2-trifluoroethanesulfonyl, and imidazolesulfonyl.
22. (Withdrawn) A method according to Claim 17, wherein L is nosylate, tosylate, or mesylate.
23. (Withdrawn) A method according to Claim 17, wherein the amine protecting group is selected from the group consisting of tert-butoxycarbamate, isopropyl carbamate, pivaloyloxymethyl carbamate, methyl carbamate, allyl carbamate, *N*-tetrahydropyran, *N*-tetrahydrofuran; t-butylamide, and *N*-pyrrolidinomethylamide.
24. (Withdrawn) A method according to Claim 17, wherein the amine protecting group is t-butoxycarbonyl.
25. (Withdrawn) A compound having the following formula:



wherein X is t-butoxycarbonyl, methyl, methoxymethyl ethyl, isobutyl, tetrahydropyranyl ether, tetrahydrofuran ether, methoxymethyl ether, bis-(2-chloroethoxy)methyl ether, 1-ethoxyethyl ether, or 1-methyl-1-methoxyethyl ether; P is an amine protecting group; and L is a leaving group.

26. (Withdrawn) A compound according to Claim 25, wherein P is selected from the group consisting of tert-butoxycarbamate, isopropyl carbamate, pivaloyloxymethyl carbamate, methyl carbamate, allyl carbamate, *N*-tetrahydropyran, *N*-tetrahydrofuran; t-butylamide, and *N*-pyrrolidinomethylamide.
27. (Withdrawn) A compound according to Claim 25, wherein P is t-butoxycarbonyl.
28. (Withdrawn) A compound according to Claim 25, wherein L is a sulfonate ester.


IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.
10/738,087	12/15/2003	Joseph C. Walsh	2003P88074US

Response To OFFICIAL ACTION

EXAMINER	
Jones, Dameron Levest	
ART UNIT	PAGE NUMBER
1618	7

29. (Withdrawn) A compound according to Claim 25, wherein L is selected from the group consisting of benzenesulfonyl, methylsulfonyl, p-methylbenzenesulfonyl, 4-nitrobenzene sulfonyl, p-bromobenzenesulfonyl, trifluoromethylsulfonyl, trichloroacetimidate, 2,2,2-trifluoroethanesulfonyl, and imidazolesulfonyl.
30. (Withdrawn) A compound according to Claim 25, wherein L is nosylate, tosylate, or mesylate
31. (Withdrawn) A compound according to Claim 25, wherein X is t-butoxycarbonyl, methyl, methoxymethyl ethyl, or isobutyl.
32. (Withdrawn) A compound having the following formula:



wherein P is an amine protecting group and L is a leaving group.


33. (Withdrawn) A compound according to Claim 32, wherein P is selected from the group consisting of tert-butoxycarbamate, isopropyl carbamate, pivaloyloxymethyl carbamate, methyl carbamate, allyl carbamate, N-tetrahydropyran, N-tetrahydrofuran; t-butylamide, and N-pyrrolidinomethylamide.
34. (Withdrawn) A compound according to Claim 32, wherein P is t-butoxycarbonyl.
35. (Withdrawn) A compound according to Claim 32, wherein L is a sulfonate ester.
36. (Withdrawn) A compound according to Claim 32, wherein L is selected from the group consisting of benzenesulfonyl, methylsulfonyl, p-methylbenzenesulfonyl, 4-nitrobenzene sulfonyl, p-bromobenzenesulfonyl, trifluoromethylsulfonyl, trichloroacetimidate, 2,2,2-trifluoroethanesulfonyl, and imidazolesulfonyl.
37. (Withdrawn) A compound according to Claim 32, wherein L is nosylate, tosylate, or mesylate.
38. (Withdrawn) A compound having the following formula:

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE


APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.
10/736,087	12/15/2003	Joseph C. Walsh	2003P88074US

Response To OFFICIAL ACTION

EXAMINER	
Jones, Dameron Levest	
ART UNIT	PAGE NUMBER
1618	8




39. (Currently amended) A method for preparing a compound having the following formula:




comprising:

a. [^{18}F]fluorinating a compound having the following formula:



wherein P is an amine protecting group and L is a leaving group, to produce a compound having the formula:

~~compound having the following formula:~~



wherein P is the same as defined above; and

b. removing the amine protecting group and Boc group to produce ^{18}F -FLT.

40. (Previously presented) A method according to Claim 39, wherein P is selected from the group consisting of tert-butoxycarbamate, isopropyl carbamate,

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.
10/736,087	12/15/2003	Joseph C. Walsh	2003P88074US

Response To OFFICIAL ACTION

EXAMINER	
Jones, Dameron Levest	
ART UNIT	PAGE NUMBER
1618	9

pivaloyloxymethyl, carbamate, methyl carbamate, allyl carbamate, *N*-tetrahydropyran, *N*-tetrahydrofuran; *t*-butylamide, and *N*-pyrrolidinomethylamide.

41. (Previously presented) A method according to Claim 39, wherein P is *t*-butoxycarbonyl.
42. (Previously presented) A method according to Claim 39, wherein L is benzenesulfonyl, methylsulfonyl, *p*-methylbenzenesulfonyl, 4-nitrobenzene sulfonyl, *p*-bromobenzenesulfonyl, trifluoromethylsulfonyl, trichloroacetimidate, 2,2,2-trifluoroethanesulfonyl, or imidazolesulfonyl.
43. (Previously presented) A method according to Claim 39, wherein L is nosylate, tosylate, or mesylate.
44. (Previously presented) A method according to Claim 39, wherein P is *t*-butoxycarbonyl and L is nosylate.
45. (Previously presented) A method according to Claim 39, wherein the amine protecting group and boc groups are removed by acid hydrolysis.
46. (Previously presented) A method according to Claim 39, wherein the amine protecting group and boc group are removed by treating the reaction product of step (a) with HCl, HBr, HOAc, H₂SO₄, HI, trimethylsilyliodide, or H₃PO₄.
47. (Withdrawn) A method for preparing a precursor for the preparation of a radiolabeled nucleoside comprising:
- converting a 2'-deoxy nucleoside into a 2,3'-anhydronucleoside;
 - opening the 2,3'-anhydro ring to produce 3'-beta-hydroxy nucleoside;
 - protecting the 5'-hydroxy with *t*-butoxycarbonyl;
 - incorporating a leaving group at the 3'-position; and
 - protecting the 3-*N* amine to produce the radiolabeled nucleoside precursor.
48. (Withdrawn) The method according to Claim 47, wherein the nucleoside is thymidine, cytidine, or uridine.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.
10/736,087	12/15/2003	Joseph C. Walsh	2003P88074US

Response To OFFICIAL ACTION

EXAMINER	
Jones, Dameron Levest	
ART UNIT	PAGE NUMBER
1618	10

49. (Withdrawn) The method according to Claim 47, wherein the leaving group is nosylate, tosylate, or mesylate.

50. (Withdrawn) The method according to Claim 47, wherein the amine protecting group is t-butoxycarbonyl.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.